

CERTIFICATE OF ANALYSIS
Botulinum Neurotoxin Type C Light Chain, Recombinant
Lot # 6253A1Contents:

Each vial of light chain from Botulinum Neurotoxin Type C contains 10 μg of lyophilized protein. When reconstituted with 120 μl of sterile distilled water, each vial contains 10 μg of light chain in 20 mM HEPES, pH 7.4 + 1.25% lactose. In order to ensure stability during storage of the protein, 0.05% TWEEN-20 or 1 mg/ml BSA must be included in the reconstitution buffer. The protein was recombinantly expressed in *E. coli* and purified using affinity and anion exchange chromatography. The affinity tag has been cleaved off of the protein prior to quantitation and packaging.

Molecular Weight:

The light chain C fragment contains amino acids 1 - 436 of the full length Botulinum Neurotoxin type C. It also contains eight residual amino acids from the affinity tag at the N - terminus to give a total length of 444 amino acids. The molecular weight of the protein has been determined to be 50, 865 Da based on electrospray-MS analysis.

Concentration:

Protein concentration was determined by a modification of the Bradford¹ method using bovine serum albumin as a standard.

Gel Electrophoresis:

When examined on 4-12% SDS-polyacrylamide gels, this product migrates as a single major band with an apparent molecular weight of approximately 50,000 Da. The protein is approximately 90% pure based on densitometry analysis.

Activity:

The recombinant light chain from Botulinum Neurotoxin Type C (LcC) has been tested for activity in a FRET based assay. Significant digestion of the FRET substrate was detected within 2 hours when using 10 and 5 nM LcC with 10 μM SYNTAXideTM substrate (Product #560).

(continued)

Packaging and Storage:

This product is supplied as a lyophilized powder which has been stoppered under vacuum. Reconstitution of the powder should be done with syringe through the rubber stopper to avoid any loss of material. Store lyophilized vials at 4°C. Once dissolved, aliquot and store at -20°C. Refrain from multiple freeze/thaw cycles.

Toxicity:

The light chain of Botulinum Neurotoxins are non-toxic and unable to penetrate cells in the absence of the heavy chain. The expression and purification of light chain from a recombinant setting ensures there is no possible contamination with heavy chain or full length intact toxin.

Handling:

Good laboratory technique should be employed in the safe handling of this product. This requires observing the following practices:

- 1. Wear appropriate laboratory attire including a lab coat, gloves and safety glasses.**
- 2. Do not mouth pipette, inhale, ingest or allow to come into contact with open wounds. Wash thoroughly any area of the body which comes into contact with the product.**
- 3. Avoid accidental autoinoculation by exercising extreme care when handling in conjunction with any injection device.**
- 4. This product is intended for research purposes by qualified personnel only. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc. is not liable for any damages resulting from the misuse or handling of this product.**

FOR RESEARCH PURPOSES ONLY. NOT FOR HUMAN USE.

References:

1. Bradford, M.M. (1976) *Anal. Biochem.* 72, 248-254.

Approved: TC Date: 7/9/09 Approved: NS Date: 7/9/07