

CERTIFICATE OF ANALYSIS
GST-SV2c, RECEPTOR FOR BOTULINUM TOXIN TYPE A
Lot # 6902A1**Contents**

Each 100 µg vial of GST-SV2c, when reconstituted with 100 µl water, contains 100 µg of protein in 20 mM HEPES, pH 7.4 + 1.25% lactose.

The protein was recombinantly expressed in *E. coli* and purified using affinity chromatography. The GST affinity tag has been retained.

Molecular Weight

GST-SV2c contains amino acids 454-579 of the full length SV2c protein. This region of the protein is known as the luminal domain loop, between transmembrane domains 7 and 8, and has been shown to be the location of botulinum neurotoxin type A (BoNT/A) binding^{1,2}. The 25 kDa GST affinity tag is on the N-terminal. The total length of the fusion protein is 330 amino acids. The molecular weight of the protein is 41,733 Da based on analysis of the nucleic acid sequence.

Concentration

Protein concentration was determined by absorbance at 280nm using Abs (0.1%) = 1.253. This value is calculated by ProtParam³ using an algorithm based on the Edelhoch⁴ method with modifications described in Pace et al⁵.

Purity

When examined on 4-12% SDS-polyacrylamide gels, this product migrates as a single major band with an apparent molecular weight of approximately 40,000 Da. The protein purity is >95% based on densitometric analysis.

Functionality

The following three posters demonstrate the functionality of the GST-SV2c fusion protein as a receptor for BoNT/A. Recent data from our laboratory indicates improved specificity and reliable detection of as little as 1.25 pg of BoNT/A when captured by the SV2c receptor. A full description of the optimized parameters will be available soon.

1. T. Christian and N. Shine. Sensitive, *In Vitro*, Bifunctional Potency Assay for Botulinum Neurotoxin Type A. The 46th Annual Interagency Botulinum Research Coordinating Committee Meeting. October, 2009 in Alexandria, VA.
2. T. Christian, A. Rummel, and N. Shine. Functional Assay for Botulinum Neurotoxin Type A Utilizing the Neuronal Receptor Protein SV2c. The 45th Annual Interagency Botulinum Research Coordinating Committee Meeting. September 2008 in Philadelphia, PA.
3. T. Christian and N. Shine. Capture Assay for Botulinum Neurotoxin Type A Utilizing the Neuronal Receptor Protein SV2c. The 6th International Conference on Basic and Therapeutic Aspects of Botulinum and Tetanus Toxins. June 2008 in Baveno, Italy.

(continued)

Packaging and Storage

This product is supplied as a lyophilized powder which has been stoppered under vacuum. Store lyophilized vials at 2-8°C. Once dissolved, aliquot and store the product at -20°C. Refrain from multiple freeze/thaw cycles.

Toxicity

GST-SV2c is non-toxic.

Handling

This product is not known to be hazardous. Good laboratory technique should be employed in the safe handling of this product. Wear appropriate laboratory attire including a lab coat, gloves and safety glasses. Nitrile gloves are recommended when handling lyophilized material.

This product is intended for research purposes only. It is not intended for use in humans or as a diagnostic agent. List Biological Laboratories, Inc. is not liable for any damages resulting from the misuse or handling of this product.

FOR RESEARCH PURPOSES ONLY. NOT FOR HUMAN USE.

References

1. Mahrhold, S., Rummel, A., Bigalke, H., Davletov, B., and Binz, T. (2006) FEBS Lett. **580**:2011-2014.
2. Dong, M., Yeh, F., Tepp, W.H., Dean, C., Johnson, E.A., Janz, R., and Chapman, E.R. (2006) Science **312**:592-596.
3. www.expasy.ch/tools/protparam-doc.html
4. Edelhoch, H. (1967) Biochemistry, **6**:1948-1954.
5. Pace, C.N., Vajdos, F., Fee, L., Grimsley, G., and Gray, T. (1995) Protein Sci., **4**:2411-2423.

Production: TC Date: 7/18/12

Management: NS Date: 7/18/12

QA/QC: QIP Date: 7/18/12